

Bibliografia

- [1] Caterson I.D., Gill T.P. Obesity epidemiology and possible prevention. *Best Pract Res Clin Endocrinol* 2002, 16, 595-610.
- [2] Prospective Studies Collaboration. Body-mass index and cause-specific mortality in 900.000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009, 373, 1083-1096.
- [3] <http://oecd.org/els/health-systems/obesityandtheconomicsofpreventionfitnotfat.htm>
- [4] Crepaldi G, Belfiore F, Bosello O, Caviezel F, Contaldo F, Enzi G, Melchionda N. Italian Consensus Conference - overweight, obesity and health. *Int J Obes* 1991 Nov; 15 (11): 781-90.
- [5] Barbone S, Castiello MR, Alborino P. *Igiene e cultura medico-sanitaria*, Franco Lucisano Editore, 2012.
- [6] Rotella CM, Ricca V, Mannucci M. *L'obesità: manuale per la diagnosi e la terapia*, SEE Firenze, 1997.
- [7] *World Health Organization Physical status: the use and interpretation of anthropometry*. Technical report series 1995, no. 854. In: Geneva: WHO, 1995.
- [8] *Obesity: The Prevention, Identification, Assessment and Management of Overweight and Obesity in Adults and Children*. Clinical guideline CG43, National Institute for Health and Clinical Excellence, London, 2006.
- [9] Greenberg JA. The obesity paradox in the US population. *Am J Clin Nutr* 2013 Jun; 97 (6): 1195-200.
- [10] Flegal KM, Graubard BI, Williamson DF, Gail MH. Excess deaths associated with underweight, overweight, and obesity. *JAMA* 2005 Apr 20; 293(15): 1861-7.
- [11] Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA* 2013 Jan 2; 309 (1): 71-82.
- [12] American Council on Exercise (ACE). What are the guidelines for percentage of body fat loss? Ask the Expert Blog. December 2, 2009.
- [13] Krakauer NY, Krakauer JC. A new body shape index predicts mortality hazard independently of body mass index. *PLoS ONE* 2012; 7(7), e39504.
- [14] Rimm EB, Stampfer MJ, Giovannucci E, et al. Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol* 1995; 141: 1117-27.
- [15] Abel Romero-Corral, Virend K. Somers, MD, Justo Sierra-Johnson, Randal J. Thomas, Kent R. Bailey, Maria L Collazo-Clavell, Thomas G. Allison, Josef Korinek, John A. Basis, Francisco Lopez-Jimenez. Accuracy of Body Mass Index to Diagnose Obesity In the US Adult Population. *Int J Obes* (Lond) 2008 June; 32 (6): 959-966.
- [16] Petursson H, Sigurdsson JA, Bengtsson C, Nilsen TIL, Getz L. Body configuration as a predictor of mortality: comparison of five anthropometric measures in a 12 year follow-up of the Norwegian HUNT 2 study. *PLoS ONE* 2001; 6: e26621.
- [17] Schoeller D.A., Jones P.J.H. Measurement of total body water by isotope dilution: a unified approach to calculations. Edited by K.J. Ellis, S. Yasamura, W.D. Morgan. In: *In vivo body composition analysis*, Institute of Physical Sciences in Medicine, London, 1987, pp. 131-137.
- [18] Wilson, Foster. Eating Disorder: obesity, anorexia nervosa, and bulimia nervosa. In: *Williams Textbook of Endocrinology*, Saunders, 1992.
- [19] Mourtzakis M, Prado CM, Liefers JR, Reiman T, McCargar LJ, Baracos VE. A practical and precise approach to quantification of body composition in cancer patients using computed tomography images acquired during routine care. *Appl Physiol Nutr Metab* 2008; 33: 997-1006.
- [20] Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Manuel Gómez J, et al. Bioelectrical impedance analysis-part II: utilization in clinical practice. *Clin Nutr* 2004; 23: 1430-53.
- [21] Tony Talluri, Sulle metodiche impedenziometriche, Akern R&D, Firenze.
- [22] Heymsfield SB, Wang ZM, Visser M, Gallagher D, Pierson RN Jr: Techniques used in the measurement of body composition: an overview with emphasis on bioelectrical impedance analysis. *Am J Clin Nutr* 1996; 64: 478S-484S.
- [23] <http://www.akern.com/en/scienceeng/theory.html>
- [24] Dicembrini I, Pala L, Cresci B, Cremasco F, Rotella CM. Predictors of weight loss in the clinical management of obese patients: the relevance of body composition. *Obes Met* 2010; 1: 29-33.
- [25] Shafer KJ, Siders WA, Johnson LK, Lukaski HC. Validity of segmental multiple-frequency bioelectrical impedance analysis to estimate body composition of adults across a range of body mass indexes. *Nutrition* 2009 Jan; 25 (1): 25-32.
- [26] Deurenberg P. Limitations of the bioelectrical impedance method for the assessment of body fat in severe obesity. *Am J Clin Nutr* 1996; 64: 449S-452S.
- [27] McCrory, M.A. et al. Evaluation of a new air displacement plethysmograph for measuring human body composition. *Med Sci Sports Exerc* 1995 Dec 27(12): 1686-91.

- [28] Jeejeebhoy K, Keith MK. Nutritional Assessment. In: Gibney M, eds. *Clin Nutr. Boston* 2005; 15-29.
- [29] Mary E. Barasi, *Human Nutrition: A Health Perspective*, CRC Press, 2003.
- [30] <http://www.orpha.net>
- [31] Kevin G. Murphy, Stephen R. Bloom. Gut hormones and the regulation of energy homeostasis. *Nature* 14 December 2006; 444, 854-859.
- [32] Kounterek SJ, Kounterek PC, Kounterek JW, Czesnikiewicz-Guzic M, Brzozowski T, Sito E. Neurohormonal Control of food intake; basic mechanism and clinical implication. *J Physiol Pharm* 2005; 56, S6: 5-25.
- [33] Tataranni PA, Gautier JE, Chen K, Uecker A, Bandy D, Salbe AD, Pratley RE, Lawson M, Reiman EM, Ravussin E. Neuroanatomical correlates of hunger and satiation in humans using positron emission tomography. *Proc Natl Acad Sci* 1999 Apr 13; 96(8): 4569-74.
- [34] Schwartz MW, Woods SC, Porte D Jr, et al. Central nervous system control of food intake. *Nature* 2000; 404: 661-671.
- [35] Führer D, Zysset S, Stumvoll M. Brain activity in hunger and satiety: an exploratory visually stimulated fMRI study. *Obesity* 2008 May; 16(5): 945-50.
- [36] McMinn JE, Baskin DG, Schwartz MW. Neuroendocrine mechanisms regulating food intake and body weight. *Obes Rev* 2000 May; 1 (1): 37-46.
- [37] Wellman PJ. Norepinephrine and the control of food intake. *Nutrition* 2000 Oct; 16 (10): 837-42.
- [38] Billington CJ, Levine AS. Hypothalamic neuropeptide Y regulation of feeding and energy metabolism. *Curr Opin Neurobiol* 1992 Dec; 2 (6): 847-51.
- [39] Kageyama H, Takenoya F, Hirako S, Wada N, Kintaka Y, Inoue S, Ota E, Ogawa T, Shioda S. Neuronal circuits involving neuropeptide Y in hypothalamic arcuate nucleus-mediated feeding regulation. *Neuropeptides* 2012 Dec; 46 (6): 285-9.
- [40] Yang K, Guan H, Arany E, Hill DJ, Cao X. Neuropeptide Y is produced in visceral adipose tissue and promotes proliferation of adipocyte precursor cells via the Y1 receptor. *FASEB J* 2008 Jul; 22 (7): 2452-64.
- [41] Bagnol D, Lu XY, Kaelin CB, Day HE, Ollmann M, Gantz I, et al. Anatomy of an endogenous antagonist: relationship between Agouti-related protein and proopiomelanocortin in brain. *J Neurosci* 1999; 19 (18): RC26.
- [42] Morton G.J., Schwartz M.W. The NPY/AgRP neuron and energy homeostasis. *Int J Obes Relat Metab Disord* 2001; 25 (Suppl. 5), S56-S62.10.1038/sj.ijo.0801915
- [43] Chen Y, Hu C, Hsu CK, Zhang Q, Bi C, Asnicar M, et al. Targeted disruption of the melanin-concentrating hormone receptor-1 results in hyperphagia and resistance to diet-induced obesity. *Endocrinology* 2003; 143(7): 2469-77.
- [44] Zigman JM, Elmquist JK. Review Minireview: From anorexia to obesity – the yin and yang of body weight control. *Endocrinology* 2003 Sep; 144 (9): 3749-56.
- [45] Pacher P, Batkai S, Kunos G. The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacol Rev* 2006; 58: 389-462.
- [46] Mechoulam R, Berry E, Avraham Y, et al. Endocannabinoids, feeding and suckling – From our perspective. *Int J Obes* 2006; 30: S24-S28.
- [47] Harrold J, Williams G. The cannabinoid system: A role in both the homeostatic and hedonic control of eating? *Br J Nutr* 2007; 90: 729-734.
- [48] Schneider JE, Buckley CA, Blum RM, Zhou D, Szymanski L, Day DE, Bartness TJ. Metabolic signals, hormones and neuropeptides involved in control of energy balance and reproductive success in hamsters. *Eur J Neurosci* 2002 Aug; 16 (3): 377-9.
- [49] Cone RD. Studies on the physiological functions of the melanocortin system. *Endocr Rev* 2006; 27 (7): 736-49.
- [50] Farooqi IS, Keogh JM, Yeo GS, Lank EJ, Cheetham T, O'Rahilly S. Clinical spectrum of obesity and mutations in the melanocortin 4 receptor gene. *N Engl J Med* 2003; 348 (12): 1085-95.
- [51] Kristensen P, Judge ME, Thim L, Ribel U, Christjansen KN, Wulff BS, et al. Hypothalamic CART is a new anorectic peptide regulated by leptin. *Nature* 1998; 393 (6680): 72-6.
- [52] Faouzi M, Leshan R, Bjornholm M, Hennessey T, Jones J, Munzberg H. Differential accessibility of circulating leptin to individual hypothalamic sites. *Endocrinology* 2007; 148 (11): 5414-23.
- [53] Cowley MA, Smart JL, Rubinstein M, Cerdan MG, Diano S, Horvath TL, et al. Leptin activates anorexigenic POMC neurons through a neural network in the arcuate nucleus. *Nature* 2001; 411 (6836): 480-4.
- [54] Baskin DG, Figlewicz Lattemann D, Seeley RJ, Woods SC, Porte D Jr, Schwartz MW. Insulin and leptin: dual adiposity signals to the brain for the regulation of food intake and body weight. *Brain Res* 1999 Nov 27; 848 (1-2): 114-23.
- [55] Sahu A. Leptin signaling in the hypothalamus: emphasis on energy homeostasis and leptin resistance. *Front Neuroendocrinol* 2003 Dec; 24 (4): 225-53.
- [56] Mayer J. Glucostatic mechanism of regulation of food intake. *N Engl J Med* 1953; 249: 13-16.
- [57] Mueckler M: Facilitative glucose transporters. *Eur J Biochem* 1994; 219: 713-725.
- [58] Air EL, Benoit SC, Blake Smith KA, Clegg DJ, Woods SC. Acute third ventricular administration of insulin decreases food intake in two paradigms. *Pharmacol Biochem Behav* 2002 May; 72 (1-2): 423-9.
- [59] Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 1999 Dec 9; 402 (6762): 656-60.
- [60] Wang Q, Liu C, Uchida A, Chuang JC, Walker A, Liu T, Osborne-Lawrence S, Mason BL, Moshier C, Berglund ED, Elmquist JK, Zigman JM. Arcuate AgRP neurons mediate orexigenic and glucoregulatory actions of ghrelin. *Mol Metab* 2013 Oct 17; 3 (1): 64-72. doi: 10.1016/j.molmet.2013.10.001. eCollection 2014.
- [61] Cummings DE, Purnell JQ, Frayo RS, Schmidova K, Wisse BE, Weigle DS. A preprandial rise in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes* 2001 Aug; 50 (8): 1714-9.
- [62] Rehfeld JF. Clinical endocrinology and metabolism. Cholecystokinin. *Best Pract Res Clin Endocrinol Metab* 2004 Dec; 18(4): 569-86.
- [63] Gibbs J, Young RC, Smith GP. Cholecystokinin decreases food intake in rats. *J Comp Physiol Psychol* 1973 84 (3): 488-95.

- [64] De Silva A, Salem V, Long CJ, Makwana A, Newbould RD, Rabiner EA, Ghatei MA, Bloom SR, Matthews PM, Beaver JD, Dhillon WS. The gut hormones PYY 3-36 and GLP-1 7-36 amide reduce food intake and modulate brain activity in appetite centers in humans. *Cell Metab* 2011 Nov 2; 14(5): 700-6.
- [65] Flint A, Raben A, Astrup A, Holst JJ. Glucagon-like peptide 1 promotes satiety and suppresses energy intake in humans. *J Clin Invest* 1998 Feb 1; 101(3): 515-20.
- [66] <http://ncbi.nlm.nih.gov/gene?Db=gene&Cmd=-ShowDetailView&TermToSearch=5697>
- [67] Alessandro Pocai. Unraveling oxyntomodulin, GLP1's enigmatic brother. *Journal of Endocrinology* 2012; 215, 335-346.
- [68] Dakin CL, Gunn I, Small CJ, Edwards CM, Hay DL, Smith DM, Ghatei MA, Bloom SR. Oxyntomodulin inhibits food intake in the rat. *Endocrinology* 2001 Oct; 142(10): 4244-50.
- [69] Rodin J, Wack J, Ferrannini E, DeFronzo RA. Effect of insulin and glucose on feeding behavior. *Metabolism* 1985 Sep; 34 (9): 826-31.
- [70] De Vos P, Saladin R, Auwerx J, Staels B. Induction of ob gene expression by corticosteroids is accompanied by body weight loss and reduced food intake. *J Biol Chem* 1995 Jul 7; 270 (27): 15958-61.
- [71] Dockray GJ. Luminal sensing in the gut: an overview. *J Physiol Pharmacol* 2003 Dec; 54 Suppl 4: 9-17.
- [72] Feldmann V, et al. UCP1 ablation induces obesity and abolishes diet-induced thermogenesis in mice exempt from thermal stress by living at thermoneutrality. *Cell Metab* 2009; 9: 203-209.
- [73] Vijgen GHEJ, Bouvy ND, Teule GJJ, Brans B, Schrauwen P, van Marken Lichtenbelt WD. Brown adipose tissue in morbidly obese subjects. *PLoS ONE* 2011; 6 (2) e17247.
- [74] *Linee Guida Italiane per l'Obesità* (LiGIO), 1999.
- [75] *Standard Italiani per la cura dell'obesità*, 2012-2013.
- [76] Serra-Majem L, Roman B, Estruch R. Scientific evidence of interventions using the Mediterranean diet: a systematic review. *Nutr Rev* 2006; 64: Suppl 1: S27-S47.
- [77] Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I, Golan R, Fraser D, Bolotin A, Vardi H, Tangi-Rozental O, Zuk-Ramot R, Sarusi B, Brickner D, Schwartz Z, Sheiner E, Marko R, Katorza E, Thiery J, Fiedler GM, Blüher M, Stumvoll M, Stampfer MJ, Dietary Intervention Randomized Controlled Trial (DIRECT) Group. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med* 2008 Jul 17; 359 (3): 229-41.
- [78] Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: meta-analysis. *BMJ* 2008 Sep 11; 337: a1344.
- [79] Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, Gómez-Gracia E, Ruiz-Gutiérrez V, Fiol M, Lapetra J, Lamuela-Raventós RM, Serra-Majem L, Pintó X, Basora J, Muñoz MA, Sorlí JV, Martínez JA, Martínez-González MA, PREDIMED Study Investigators. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013 Apr 4; 368(14): 1279-90.
- [80] Lagiou P, Sandin S, Lof M, Trichopoulos D, Adami HO, Weiderpass E. Low carbohydrate-high protein diet and incidence of cardiovascular diseases in Swedish women: prospective cohort study. *BMJ* 2012 Jun 26; 344: e4026.
- [81] Lars Osterberg, Terrence Blaschke. Drug Therapy – Adherence to Medication. *N Eng J Med* 2005; 353: 487-497.
- [82] DiMatteo M., *Med Care* 2004; 42: 200-290.
- [83] Bier DM et al. Improving Compliance with Dietary Recommendations: Time for New, Inventive Approaches? *Nutrition Today* 2008; Vol. 43 No. 5, 180-187.
- [84] Masley S., *The Permanente Journal/Summer* 1998; Vol. 2 No. 3, 12-16.
- [85] S.H.A. Holt, J.C. Brand Miller, P. Petocz, E. Farmakalidis, A Satiety Index of Common Foods, *European Journal of Clinical Nutrition*, September 1995, 675-690.
- [86] Alan W. Barclay, Jennie Brand-Miller. The Australian Paradox: A Substantial Decline in Sugars Intake over the Same Timeframe that Overweight and Obesity Have Increased. *Nutrients* 2011; 3 (4), 491-504.
- [87] Björvell H, Rossner S. A ten-year follow-up of weight change in severely obese subjects treated in a combined behavioural modification programme. *Int J Obes Relat Metab Disord* 1992; 16: 623-625.
- [88] Wadden TA, Foster GD. Behavioral treatment of obesity. *Med Clin North Am* 2000; 84: 441-461.
- [89] Alisivatos JG. Patient Education: individual, group, and mass media approach. Advantages and disadvantages. In: Assal JP, Berger M, Gay N, Carnivet J (eds). *Diabetes Education. How to improve patients' education*, Excerpta-Medica, Amsterdam, 1983, 130-138.
- [90] Mannucci E, Conti A, Rotella CM. Gruppo interattivo di educazione e di sostegno (GIES): un nuovo approccio all'educazione permanente nei diabetici tipo 1. *G It Diabetol* 1994; 14: 243.
- [91] Verweij LM, Coffeng J, van Mechelen W, Proper KJ. Meta-analyses of workplace physical activity and dietary behaviour interventions on weight outcomes. *Obes Rev* 2011; 12 (6): 406-29.
- [92] B. Cresci, F. Tesi, T. La Ferlita, V. Ricca, C. Ravaldi, C.M. Rotella, E. Mannucci. Group versus individual cognitive-behavioral treatment for obesity: results after 36 months. *Eat Weight Disord* 2007; 12 (4): 147-53.
- [93] Minniti A, Bissoli L, Di Francesco V, Fantin F, Mandragona M, Olivieri M, Fontana G, Rinaldi C, Bosello O, Zamboni M. Individual versus group therapy for obesity: comparison of drop-out rate and treatment outcome. *Eat Weight Disord* 2007; 12 (4): 161-7.
- [94] Cresci B, Castellini G, Pala L, Bigiarini M, Romoli E, Poggiali R, Guarnieri C, Biffi B, La Ferlita T, Ricca V, Mannucci E, Rotella CM. Fit and motivated: outcome predictors in patients starting a program for lifestyle change. *Obes Facts* 2013; 6 (3): 279-87.
- [95] Jeffrey RW, Forster JL, French SA, et al. The Healthy Worker Project: a worksite intervention for weight control and smoking cessation. *Am J Pub Health* 1993; 89: 325.
- [96] Perri MG, McAdu WG, McAllister DA et al. Effect of peer support and therapist control on long term weight loss. *J Consult Clin Psychol* 1987; 55: 615-617.
- [97] Avila P, Hovell MF. Physical activity and training for weight loss in Latinas: a controlled trial. *Int J Obes Relat Metab Disorder* 1994; 18: 476.

- [98] Cresci B, Rotella CM. Motivational readiness to change in lifestyle modification programs. *Eat Weight Disord* 2009.
- [99] Cresci B, Castellini G, Pala L, Ravaldi C, Faravelli C, Rotella CM, Ricca V. Motivational readiness for treatment in weight control programs: the TREATment Motivation and REadiness (TRE-MORE) test. *J Endocrinol Invest* 2011 Mar; 34 (3): e70-7.