

Aspetti dietetici nelle malattie renali

Bibliografia

- [1] Abboud H, Henrich WL. Clinical practice. Stage IV chronic kidney disease. *N Engl J Med* 2010; 362: 56-65.
- [2] Acchiardo AR, Moore LW, Latour PA. Malnutrition as the main factor in morbidity and mortality of hemodialysis patients. *Kidney Int* 1983; 24: S199-S203.
- [3] Aparicio M, Bellizzi V, Chauveau P, Cupisti A, et al. Do ketoanalogues still have a role in delaying dialysis initiation in CKD predialysis patients? *Semin Dial* 2013; 26 (6): 714-719.
- [4] Aparicio M, Bellizzi V, Chauveau P, Cupisti A, et al. Keto acid therapy in predialysis chronic kidney disease patients: final consensus. *J Ren Nutr* 2012; 22 (2Supp): S22-4.
- [5] Aparicio M, Chauveau P, Combe C. Low protein diets and outcome of renal patients. *J Nephrol* 2001; 14: 433-439.
- [6] Ash S, Campbell KL, Bogard J, Millichamp A. Nutrition prescription to achieve positive outcomes in chronic kidney disease: a systematic review. *Nutrition* 2014; 6: 416-451.
- [7] Attman PO, Samuelsson O. Dyslipidemia of kidney disease. *Curr Opin Lipidol* 2009; 20: 293-9.
- [8] Australian Institute of Health and Welfare. *Projections of the Incidence of Treated End-Stage Kidney Disease in Australia*, AIHW, Canberra, 2011.
- [9] Barsotti G. Historical archives of Italian nephrology: Sergio Giovanetti (1924-2000): a distinguished representative of the school of Pisa. *G Ital Nefrol* 2006; 23 (3): 350-7.
- [10] Barsotti G, Morelli E, Cupisti A, et al. A special supplemented «vegetarian» diet for nephrotic patient. *Am J Nephrol* 1991; 11: 380-385.
- [11] Beale LS. *Kidney disease, urinary deposits and calculous disorders: their nature and treatment*. Philadelphia, Lindsay and Blakiston, 1869.
- [12] Bernstein AM, Treyzon L, Li Z. Are high-protein, vegetable-based diet safe for kidney function? A review of the literature. *J Am Diet Assoc* 2007; 107: 644-50.
- [13] Bossola M, Muscaritoli M, Tazza L, et al. Malnutrition in hemodialysis patients: what therapy? *Am J Kidney Dis* 2005; 46: 371-86.
- [14] Brunini TMC, Roberts NB, Yaqoob MM, Ellory JC, et al. Activation of L-arginine transport in undialysed chronic renal failure and continuous ambulatory peritoneal dialysis patients. *Clin Exp Pharmacol Physiol* 2006; 33 (12): 114-118.
- [15] Brunori G. Nutrition in chronic renal disease and renal failure. In: Bales CW and Ritchie CS (eds). *Handbook of clinical nutrition and aging*, Humana Press, 2003, pp. 599-6219.
- [16] Brunori G, Viola BF, Parrinello G, et al. Efficacy and safety of a very-low-protein diet when postponing dialysis in the elderly: a prospective randomized multi-center controlled study. *Am J Kidney Dis* 2007; 49: 569-80.
- [17] Canepa A, Filho JC, Gutierrez A, et al. Free amino acids in plasma, red blood cells, polymorphonuclear leukocytes, and muscle in normal and uraemic children. *Nephrol Dial Transplant* 2002; 17 (3): 413-421.
- [18] Carrero JJ, Qureshi AR, Axelsson J, et al. Comparison of nutritional and inflammatory markers in dialysis patients with reduced appetite. *Am J Clin Nutr* 2007; 85: 695-701.
- [19] Castellino P. Protein metabolism in renal failure. *G Ital Nefrol* 2005; 22: 581-589.
- [20] Chauveau P, Couzi L, Vendrely B, de Precigout V, Combe C, Fouque D, Aparicio M. Long-term outcome on renal replacement therapy in patients who previously received a keto acid-supplemented very-low-protein diet. *Am J Clin Nutr* 2009; 90, 969-974.
- [21] Cianciaruso B, Barsotti G, Del Vecchio L, et al. Linee Guida per la terapia conservativa della insufficienza renale cronica. *G Ital Nefrol* 2003; 20: 48-60.
- [22] Cianciaruso B, Bellizzi V, Minutolo R, et al. Salt intake and renal outcome in patients with progressive renal disease. *Miner Electrolyte Metab* 1989; 24: 272-279.
- [23] Cianciaruso B, Brunori G., *Terapia nutrizionale nel paziente con insufficienza renale cronica*. Collana Aggiornamenti in tema di nefrologia, Forum Service Editore, 2005.
- [24] Cupisti A, D'Alessandro C, Morelli E et al. Nutritional status and dietary manipulation in predialysis chronic renal failure patients. *J Ren Nutr* 2004; 14: 127-133.
- [25] Curhan, GC, Willett WC, Rimm EB, Stampfer MJ. A Prospective Study of Dietary Calcium and Other Nutrients and the Risk of Symptomatic Kidney Stones. *N Engl J Med* 1993; 328: 833-838.
- [26] Curhan, GC, Willett WC, Knight EL et al. Dietary factors and the risk of incident kidney stones in younger women: nurses' wealth study II. *Arch Intern Med* 2004 Apr 26; 164 (8): 885-91.
- [27] Dumont, R., Lloyd, L., Meade, A. et al. Evidence based practice guidelines for the nutritional management of chronic kidney disease. *Nutr Diet* 2006; 63, S35-S45.
- [28] Dussol B, Iovanna C, Raccach D et al. A randomized trial of low-protein diet in type 1 and type 2 diabetes mel-

- litus patients with incipient and overt nephropathy. *J Ren Nutr* 2005; 15: 398-406.
- [29] El Haggan W, Vendrely B, Chauveau P et al. Early evolution of nutritional status and body composition after kidney transplantation. *Am J Kidney Dis* 2002; 40: 629-637.
- [30] FAO/WHO/UNO. Energy and protein requirements. In: *Technical Report Series. World Health Organization* 1985; 724: 1-206.
- [31] Fatina I, Fadel, Manal F, Elshamaa, Rascha G, Essam, Eman A., Elghoroury Gamila S. M., El-Saeed Safinaz E., El-Toukhy, Mona Hamed Ibrahim. Some Amino Acids Levels: Glutamine, Glutamate, and Homocysteine, in Plasma of Children with Chronic Kidney Disease. *Int J Biomed Sci* 2014 Mar; 10 (1): 36-42.
- [32] Fogazzi G.B. Archivi storici della nefrologia italiana: Gabriele Monasterio (1903-1972) e la scuola pisana. *G Ital Nefrol* 2005; 6: 597-608.
- [33] Fouque D, Laville M, Boissel JP. Low protein diets for chronic kidney disease in non diabetic adults. *Cochrane Database Syst Rev* 2009; 19: CD001892.
- [34] Garibotto G, Sofia A, Saffiotti S, Bonanni A, Mannucci I, Parodi EL, Cademartori V, Verzola D. Effects of peritoneal dialysis on protein metabolism. *Nutr Metab Cardiovasc Dis* 2013; 23 Suppl 1: S25-30.
- [35] Gifford JD, Rustky EA, Kirk KA et al. Control of serum potassium during fasting in patients with end stage renal disease. *Kidney Int* 1989; 35: 90-94.
- [36] Giovannetti S. Low-protein diets for chronic renal failure. In: Giovannetti S (ed), *Nutritional treatment of chronic renal failure*, Kluwer Academic Publishers, 1989, 179-190.
- [37] Giovannetti S, Maggiore Q. A low-nitrogen diet with proteins of high biological value for severe chronic uremia. *Lancet* 1964; 37: 1000-1003.
- [38] Hulley S, Grady D, Bush T, et al. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and Estrogen/progestin Replacement Study (HERS) Research Group. *JAMA* 1998; 280: 605-613.
- [39] Johnson, D.W., Atai, E., Chan, M., Phoon, R.K.S., Scott, C.; Toussaint, N.D., Turner, G.L., Usherwood, T., Wiggins, K.J. KHA-CARI guideline: Early chronic kidney disease: Detection, prevention and management. *Nephrology* 2013; 18, 340-350.
- [40] Kasiske BL. Hyperlipidemia in patients with chronic renal disease. *Am J Kidney Dis* 1998; 32: S142-56.
- [41] Kasiske BL, Lakatua JD, Ma JZ et al. A meta-analysis of the effects of dietary protein restriction on the rate of decline in renal function. *Am J Kidney Dis* 1998; 31: 954-961.
- [42] Kaysen GA. Lipid and lipoprotein metabolism in chronic kidney disease. *J Ren Nutr* 2009; 19: 73-7.
- [43] Klahr S, Levey AS, Beck GJ et al. The effects of dietary protein restriction and blood pressure control on progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. *N Engl J Med* 1994; 330: 877-884.
- [44] KDIGO Kidney Disease: Improving Global Outcomes CKD Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int* 2013; Suppl. 3, 1-150.
- [45] K/DOQI clinical practice guidelines for bone metabolism and disease in chronic kidney disease. *Am J Kidney Dis* 2003; 42: 1-201.
- [46] Kopple JD. Pathophysiology of protein-energy wasting in chronic renal failure. *J Nutr* 1999; 129 (1S Suppl): 247S-251S.
- [47] Krenitsky J. Nutrition in renal failure: myths and management. *Pract gastroenterol* 2004; 9: 40-59.
- [48] Kwan BC, Kronenberg F, Beddhu S, Cheung AK. Lipoprotein metabolism and lipid management in chronic kidney disease. *J Am Soc Nephrol* 2007; 18: 1246-61.
- [49] Levey AS, Adler S, Caggiula AW et al. Effects of dietary protein restriction on progression of advanced renal disease in the Modification of Diet in Renal Disease study. *Am J Kidney Dis* 1996; 27: 652-663.
- [50] Levey AS, Greene T, Beck GJ et al. Dietary Protein Restriction and the Progression of Chronic Renal Disease: What Have All of the Results of MDRD Study Shown? *J Am Soc Nephrol* 1999; 10: 2426-2439.
- [51] Levey AS, Greene T, Sarnak MJ et al. Effect of dietary protein restriction on progression of kidney disease: long-term follow-up of the Modification of Diet in Renal Disease (MDRD) study. *Am J Kidney Dis* 2006; 48: 879-888.
- [52] Levey AS, Lesley AS, Schmid CH, Zhang YL, Castro A, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J for the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI). *Ann Intern Med* 2009; 150 (9): 604-61.
- [53] Lim VS, Kopple JD. Protein metabolism in patients with chronic renal failure: role of uremia and dialysis. *Kidney Int* 2000; 58 (1): 1-10.
- [54] *Linee guida nazionali: identificazione, prevenzione e gestione della malattia renale cronica dell'adulto*, ISS, 2012.
- [55] Linee guida per la terapia conservativa della insufficienza renale cronica. *GIN* 2003; S24: S48-S60.
- [56] Locatelli F, Alberti D, Graziani G et al. Prospective, randomized multicentre trial of effect of protein restriction on progression of chronic renal insufficiency. Northern Italian cooperative Study Group. *Lancet* 1991; 337: 1299-1304.
- [57] Maeda Y, Shiigai T. Diet in diabetic nephropathy. *Contrib Nephrol* 2007; 155: 50-58.
- [58] Maschio G, Oldrizzi L, Tessitore N et al. Effects of dietary protein and phosphorus restriction on progression of early renal failure. *Kidney Int* 1982; 22: 371-376.
- [59] Mehrotra R, Kopple JD. Nutritional management of maintenance dialysis patients: why aren't we doing better? *Ann Rev Nutr* 2001; 21: 343-379.
- [60] Mitch WE. Chronic kidney disease. In: Goldman L, Ausiello D, eds., *Cecil Medicine*, 23rd ed., Saunders Elsevier, 2007, chap. 131.
- [61] Mitch WE. Requirements for protein, calories and fat in the predialysis patient. In *Handbook of Nutrition and the Kidney*. Mitch WE and Klahr S (eds), Philadelphia, Lippincott, Williams and Wilkins: 144-165, 2002.
- [62] Mitch WE, Remuzzi G. Diets for patients with chronic kidney disease, still worth prescribing. *J Am Soc Nephrol* 2004; 15: 234-237.
- [63] Magorzewicz S, Debska-Slizien A, Rutkowski B, Lysiak-Szydowska W. Serum concentration of amino acids

- versus nutritional status in hemodialysis patients. *J Ren Nutr* 2008; 18 (2): 239-247.
- [64] Moist LM, Port FK, Orzol SM et al. Predictors of loss of residual renal function among new dialysis patients. *J Am Soc Nephrol* 2000; 11: 556-564.
- [65] Molino A, Chiappini MG, Laviano A, Ammann T, Bollea MR, Alegiani F, Rossi Fanelli F, Muscaritoli M. Effect of intensive nutritional counseling and support on clinical outcomes of hemodialysis patients. *Nutrition* 2012; 28 (10): 1012-5.
- [66] Moreau K, Chauveau P, Martin S et al. Long-term evolution of body composition after renal transplantation: 5-year survey. *J Ren Nutr* 2006; 16: 291-299.
- [67] Muscaritoli M, Molino A, Bollea MR, Fanelli F, Malnutrition and wasting in renal disease. *Curr Opin Clin Nutr Metab Care* 2009; 12: 378-383.
- [68] National Kidney Foundation: DOQI Clinical Practice Guidelines for Hemodialysis and Peritoneal Dialysis Adequacy. *Am J Kidney Dis* 1997; 30: 1-136.
- [69] National Kidney Foundation. *Dietary Guidelines for Adults Starting on Hemodialysis*, 2009.
- [70] Navaneethan, S.D., Yehnert, H., Moustarah, F., Schreiber, M.J., Schauer, P.R., Beddhu, S. Weight loss interventions in chronic kidney disease: A systematic review and meta-analysis. *Clin J Am Soc Nephrol* 2009; 4, 1565-1574.
- [71] Nishi S, Gejyo F, Saito K et al. Diet therapy after kidney transplantation. In: Suzuki H, Kimmel PL (eds), *Nutrition and kidney disease: a new era*. Contrib Nephrol vol 155, Karger, 2007, 82-89.
- [72] Obrador GT, Arora P, Kausz AT et al. Level of renal function at the initiation of dialysis in the US end-stage renal disease population. *Kidney Int* 1999; 56: 2227-2235.
- [73] Pecoits-Filho R. Dietary protein intake and kidney disease in western diet. *Contrib Nephrol* 2007; 155: 102-112.
- [74] Piccoli GB, Motta D, Martina G et al. Low-protein vegetarian diet with alpha-chetoanalogues prior to pre-emptive pancreas-kidney transplantation. *Rev Diabet Stud* 2004; 1 (2): 95-102.
- [75] Pupim LB, Cuppari L, Ikizler TA. Nutrition and metabolism in kidney disease. *Semin Nephrol* 2006; 26: 134-157.
- [76] Robertson, L.M., Waugh, N., Robertson, A. Protein restriction for diabetic renal disease. *Cochrane Database Syst Rev* 2007; CD002181.
- [77] Strid H, Simren M, Stotzer P et al. Patients with chronic renal failure have abnormal small intestinal motility and a high prevalence of small intestinal bacterial overgrowth. *Digestion* 2003; 67: 129-137.
- [78] United States Renal Data System: 2002 annual report. Incidence and prevalence of ESRD. *Am J Kidney Dis* 2003; 42: 1-230.
- [79] Voormolen N, Noordzij M, Grootendorst DC et al. High plasma phosphate as a risk factor for decline in renal function and mortality in predialysis patients. *Nephrol Dial Transplant* 2007; 22: 2929-16.
- [80] Weir MR. Is it the low-protein diet or simply the salt restriction? *Kidney Int* 2007; 71: 188-190.
- [81] Xu H, Huang X, Arnlov J, Cederholm T et al. Clinical correlates of insulin sensitivity and its association with mortality among men with CKD stages 3 and 4. *Clin J Am Soc Nephrol* 2014; 9 (4): 690-7.
- [82] Greene T. What did we learn from the HEMO Study? Implications of secondary analyses. *Contrib Nephrol* 2005; 149-82.
- [83] Frassetto LA, Lanham-New SA, Macdonald HM, Remer T, Sebastian A, Tucker KL, Tyllavsky FA, Standardizing terminology for estimating the diet-dependent net acid load to the metabolic system. *J Nutr* 2007; 137: 1491-1492.
- [84] Ketteler M. Phosphate Metabolism in CKD Stages 3-5: Dietary and Pharmacological Control. *Int J Nephrol* 2011: Article ID 970245, 6 pages.