

Bibliografia

- [1] Mancia G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Böhm M, Christiaens T, Cifkova R, De Backer G, Dominiczak A, Galderisi M, Grobbee DE, Jaarsma T, Kirchhof P, Kjeldsen SE, Laurent S, Manolis AJ, Nilsson PM, Ruilope LM, Schmieder RE, Sirnes PA, Sleight P, Viigimaa M, Waeber B, Zannad F, Task Force Members. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2013; 31: 1281-357.
- [2] Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380 (9859): 2224-2260.
- [3] <http://www.dablededucational.org>
- [4] Nishida C, Uauy R, Kumanyika S, Shetty P. The joint WHO/FAO expert consultation on diet, nutrition and the prevention of chronic diseases: process, product and policy implications. *Public Health Nutr* 2004; 7 (1A): 245-50.
- [5] Núñez-Córdoba JM1, Valencia-Serrano F, Toledo E, Alonso A, Martínez-González MA. The Mediterranean diet and incidence of hypertension: the Seguimiento Universidad de Navarra (SUN) Study. *Am J Epidemiol* 2009 Feb 1; 169(3): 339-46.
- [6] Toledo E, de A Carmona-Torre F, Alonso A, Puchau B, Zulet MA, Martínez JA, Martínez-González MA. Hypothesis-oriented food patterns and incidence of hypertension: 6-year follow-up of the SUN (Seguimiento Universidad de Navarra) prospective cohort. *Public Health Nutr* 2010 Mar; 13 (3): 338-49.
- [7] <http://openheart.bmj.com/content/1/1/e000167.full>
- [8] Aburto NJ, Ziolkovska A, Hooper L, Elliott P, Cappuccio FP, Meerpohl JJ. Effect of lower sodium intake on health: systematic review and meta-analyses. *BMJ* 2013 Apr 3; 346: f1326.
- [9] Mendis S, Lindholm LH, Mancia G, Whitworth J, Alderman M, Lim S, Heagerty T. World Health Organization (WHO) and International Society of Hypertension (ISH) risk prediction charts: assessment of cardiovascular risk for prevention and control of cardiovascular disease in low and middle-income countries. *J Hypertens* 2007; 25: 1578-1582.
- [10] WHO. Guideline: Sodium intake for adults and children. Geneva, World Health Organization (WHO), 2012.
- [11] George J, Majeed W, Mackenzie IS, MacDonald TM, Wei L. Association between cardiovascular events and sodium-containing effervescent, dispersible, and soluble drugs: nested case-control study. *BMJ* 2013; 347: f6954.
- [12] Aburto NJ, Hanson S, Gutierrez H, Hooper L, Elliott P, Cappuccio FP. Effect of increased potassium intake on cardiovascular risk factors and disease: systematic review and meta-analyses. *BMJ* 2013; 346: f1378.
- [13] Xin X, He J, Frontini MG, Ogden LG, Motsamai OI, Whelton PK. Effects of alcohol reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 2001; 38: 1112-7.
- [14] Galletti F, Agabiti-Rosei E, Bernini G, Boero R, Desideri G, Fallo F, Mallamaci F, Morganti A, Castellano M, Nazzaro P, Trimarco B, Strazzullo P, MINISAL-GIRCSI Program Study Group. Excess dietary sodium and inadequate potassium intake by hypertensive patients in Italy: results of the MINISAL-SIIA study program. *J Hypertens* 2014; 32: 48-56.
- [15] Noyan Gokce. L-Arginine and Hypertension. *J Nutr* October 1, 2004; vol. 134, no. 10, 2807S-2811S.
- [16] Mehta S, Stewart DJ, Levy RD. The hypotensive effect of L-arginine is associated with increased expired nitric oxide in humans. *Chest* 1996 Jun; 109 (6): 1550-5.