

## Bibliografia

- [1] Chernoff R. An Overview of Tube Feeding: From Ancient Times to the Future. *Nutrition in Clinical Practice* 2006; 21: 408-410.
- [2] Dudrick SJ, Wilmore DW, Vars HM, Rhoads JE. Long-term parenteral nutrition with growth, development, and positive nitrogen balance. *Surgery* 1968; 64: 134-142.
- [3] Driscoll DF, Bistrrian BR. Parenteral and Enteral Nutrition in the Intensive Care Unit. In: *Irwin and Rippe's Intensive Care Medicine*, Lippincott Williams & Wilkins, 2008.
- [4] American Society for Parenteral and Enteral Nutrition Board of Directors. Definition of terms used in A.S.P.E.N. guidelines and standards. *JPEN* 1995; 19: 1-2.
- [5] Robinson G, Goldstein M, Levine GM. Impact of nutritional status on DRG length of stay. *JPEN* 1987; 11: 49-52.
- [6] Edington J, Kon P, Martyn CN. Prevalence of malnutrition in patients in general practice. *Clin Nutr* 1996; 15: 60-3.
- [7] Braga M, Baccani P, Scarabarozzi S et al. Prognostic role of preoperative nutritional and immunological assessment in the surgical patients. *JPEN* 1988; 12 (2): 138-42.
- [8] Leiter LA, Marliss EB. Survival during fasting may depend on fat as well as protein stores. *JAMA* 1982; 248: 2306-7.
- [9] Candusso M, Braga M, Faraguna D et al. Fogli di Trieste. *RINPE* 1994; 12: 107-10.
- [10] Schiesser M, Mueller S, Kirchoff P, Bretenstein S, Schaefer M, Clavien PA. Assessment of a novel screening score for nutritional risk in predicting complications in gastro-intestinal surgery. *Clin Nutr* 2008; 27: 565-70.
- [11] Suchner U, Kuhn KS, Furst P. The scientific basis of immunonutrition. *Proc Nutr Soc* 2000; 59: 553-63.
- [12] Heys SD, Walker LG, Smith I, Eremin O. Enteral nutrition supplementation with key nutrients in patients with critical illness and cancer. A meta-analysis of randomized controlled clinical trial. *Ann Surg* 1999; 229: 467-77.
- [13] Beale RJ, Bryg DJ, Bihari DJ. Immunonutrition in the critically ill: a systematic review on clinical outcome. *Crit Care Med* 1999; 27: 2799-8.
- [14] Wilmore D, Shabert JK. Role of glutamine in immunological responses. *Nutrition* 1998; 14: 618-26.
- [15] Drover JW, Dhaliwal R, Wischmeyer PE, Ochoa JB, Heyland DK. Perioperative use of arginine-supplemented diets: a systematic review of the evidence. *J Am Coll Surg* 2011; 212: 385-99.
- [16] Lieberman JM, Goldfine J, Shou J et al. Enteral nutrition with supplemental arginine, RNA, and omega-3 fatty acids in patients after operation: immunologic, metabolic and clinical outcome. *Surgery* 1992; 112: 56-67.
- [17] Efron D, Barbul A. Role of arginine in immunonutrition. *J Gastroenterol* 2000; 35 Suppl XII: 20-3.
- [18] Redmon HP, Stapleton PP, Neary P et al. Immunonutrition: the role of taurine. *Nutrition* 1998; 14: 599-604.
- [19] Charney P. Nutrition Assessment in the 1990's, where are we now? *Nutr Clin Pract* 1995; 10: 131-9.
- [20] Mears E. Outcomes of continuous process improvement of a nutritional care program incorporating serum prealbumin measurements. *Nutrition* 1996; 12: 479-84.
- [21] Società Italiana di Nutrizione Umana, LARN - Livelli di Assunzione Raccomandati di energia e Nutrienti per la popolazione italiana, Istituto Nazionale della Nutrizione e Ministero dell'Agricoltura e Foreste, 1996.
- [22] WHO, *Energy and Protein requirements: Report of a joint FAO/WHO/UNU expert consultation*, WHO Technical Report Series 724, 1985.
- [23] National Research Council, *Recommended Dietary Allowances*, 10th ed., Academy Press, 1998.
- [24] A.S.P.E.N. Board of Directors and the Clinical Guidelines Task Force. Guidelines for the use of Parenteral and Enteral Nutrition in adult and pediatric patients. *JPEN* 2002; 26 (Suppl 1): 52SA.
- [25] National Advisory Group on Standards and Practice Guidelines for Parenteral Nutrition. Safe practices for parenteral nutrition formulations. *JPEN* 1998; 22: 49-66.
- [26] Kris-Etherton PM, Taylor DS, Yu-Poth S et al. Polysaturated fatty acids in the food chain in the United States. *Am J Clin Nutr* 2000; 71 (suppl): S179-88.
- [27] Orr ME. Vascular access device selection for parenteral nutrition. *Nutr Clin Pract* 1999; 14: 172-7.
- [28] Van den Berghe, Wilmer A, Hermans G, Meersman W et al. Intensive insulin therapy in the medical ICU. *N Engl J Med* 2006; 354 (5): 449-61.
- [29] NICE-SUGAR Study Investigators, Finfer S, Chittock DR, Su SY, Blair D et al. Intensive versus conventional glucose control in critically ill patients. *N Eng J Med* 2009 Mar 26; 360 (13): 1283-97.
- [30] Staun M, Pironi L, Bozzetti F, Baxter J et al. ESPEN Guidelines on Parenteral Nutrition: Home Parenteral Nutrition (HN) in adult patients. *Clinical Nutrition* 2009 (28): 467-479.

- [31] Singer P, Berger MM, Van den Berghe G, Biolo G et al. ESPEN Guidelines on Parenteral Nutrition: Intensive Care. *Clinical Nutrition* 2009 (28): 387-400.
- [32] A.S.P.E.N Clinical Guidelines: Parenteral Nutrition Ordering, Order Review, Compounding, Labeling, and Dispensing. *JPEN* 2014; 38 (3): 334-377.
- [33] Rose W. The amino acid requirements of adult man. *Nutr Abstr Rev* 1957; 27: 631-47.
- [34] Frankel W, Zhang W, Singh A et al. Fiber: Effect on bacterial translocation and intestinal mucin content. *World J of Surg* 1995; 19: 144-148.
- [35] Hale Akbaylar. Basic principles of enteral nutrition. *Turk J Gastroenterol* 2002; 13(4): 186-191.
- [36] Gaggiotti G., *Manuale di Nutrizione Enterale*, Minerva Medica, 1990.
- [37] Matuchansky C. Influence de la voie de nutrition, entérale ou parentérale, sur l'immunité, la translocation, la perméabilité et la trophicité intestinales. *Nutr Clin Métabol* 1994; 8: 135-148.
- [38] Mullen JL, Buzby GP, Waldman MT et al. Prediction of operative morbidity and mortality by preoperative nutritional assessment. *Surg Forum* 1979; 30: 80-2.
- [39] Cuthbertson DP. Post-shock metabolic response. *Lancet* 1942; vol. 239, no. 6189, 433-436.
- [40] Kehlet H, Wilmore DW. Multimodal strategies to improve surgical outcome. *The American Journal of Surgery* 2002; 183: 630-641.
- [41] Macias J, Alaka KJ, Murphy MH et al. Impact of the nutritional regimen on protein catabolism and nitrogen balance in patients with renal failure. *JPEN* 1996; 20: 56-62.
- [42] Fiaccadori E, Lombardi M, Leonardi S et al. Prevalence and clinical outcome associated with preexisting malnutrition in acute renal failure: a prospective cohort study. *J Am Soc Nephrol* 1999; 10: 581-93.
- [43] Kopple JD. The nutrition management of the patients with acute renal failure. *JPEN* 1996; 20 (1): 98-107.
- [44] Simpson F, Doig GS. Parenteral vs. enteral nutrition in the critically ill patient: a meta-analysis of trial using the intention to treat principle. *Int Care Med* 2005; 31: 12-23.
- [45] Kreymann KG, Berger MM, Deutz NE et al. ESPEN Guidelines on enteral nutrition: intensive care. *Clin Nutr* 2006; 25: 210-23.
- [46] Heyland DK, Schroter-Noppe D, Drover JW et al. Nutritional support in the critical care setting: current practice in Canadian ICUs – opportunities for improvement? *JPEN* 2003; 27: 74-83.
- [47] Marik PE, Zaloga GP. Early enteral nutrition in acutely ill patients: a systematic review. *Crit Care Med* 2001; 29: 2264-70.
- [48] Muller MJ. Malnutrition in cirrhosis. *J Hepatol* 1995; 23 (Suppl 1): 33.
- [49] Italian Multicentre Cooperative Project on nutrition in liver cirrhosis. Nutritional status in cirrhosis. *J Hepatol* 1994; 21: 317-25.
- [50] Dolz C, Raurich JM, Ibáñez J, Obrador A, Marsè P, Gay J. Ascites increased the resting energy expenditure in liver cirrhosis. *Gastroenterology* 1991; 100: 738-44.
- [51] Bianchi G, Marchesini G, Zoli M, Bugianesi E, Fabbrì A, Pisi E. Prognostic significance of diabetes in patient with cirrhosis. *Hepatology* 1994; 20: 119-25.
- [52] Burke PA, Ling PR, Forse RA, Lewis DW, Jenkins R, Bistrrian BR. Sites of conditional essential fatty acid deficiency in end stage liver disease. *JPEN* 2001; 25 (4): 188-93.
- [53] Greenberg GR, Fleming CR, Jeejeebhoy KN, Rosenberg IH, Sales D, Tremaine WJ. Controlled trial of bowel rest and nutritional support in the management of Crohn's disease. *Gut* 1988; 29: 1309-15.
- [54] Gonzàles-Huix F, Fernandez-Banares F, Esteve-Comas M et al. Enteral versus parenteral nutrition as adjunct therapy in acute ulcerative colitis. *Am J Gastroenterol* 1993; 88: 227-32.
- [55] Christie PM, Hill GL. Effect of intravenous nutrition on nutrition and function in acute attacks of inflammatory bowel disease. *Gastroenterology* 1990; 99: 730-6.
- [56] Steinhart AH, Hiruki T, Brzezinski A et al. Treatment of left-sided ulcerative colitis with Butyrate enemas: a controlled trial. *Aliment Pharmacol Ther* 1996; 10: 729-36.
- [57] Scheppach W, Muller JG, Boxberger F et al. Histological changes in the colonic mucosa following irrigation with short-chain fatty acids. *Eur J Gastroenterol Hepatol* 1997; 9: 149-53.
- [58] Breuer RI, Soergel KH, Lashner BA et al. Short chain fatty acid rectal irrigation for left-sided ulcerative colitis: a randomised, placebo controlled trial. *Gut* 1997; 40: 485-91.
- [59] Vernia P, Monteleone G, Grandinetti G et al. Combined oral sodium butyrate and mesalazine treatment compared to oral mesalazine alone in ulcerative colitis: randomised, double-blind, placebo-controlled pilot study. *Dig Dis Sci* 2000; 45: 976-81.
- [60] Aslan A, Triadafilopoulos G. Fish oil fatty acid supplementation in active ulcerative colitis: a double blind, placebo-controlled, crossover study. *Am J Gastroenterol* 1992; 87: 423-7.
- [61] Hawthorne AB, Daneshmend TK, Hawkey CJ et al. Treatment of ulcerative colitis with fish oil supplementation: a prospective 12 month randomised controlled trial. *Gut* 1992; 33: 922-8.
- [62] Lorenz R, Weber PC, Szimnau P et al. Supplementation with n-3 fatty acids from fish oil in chronic inflammatory bowel disease – a randomized, placebo-controlled, double-blind cross-over trial. *J Intern Med* 1989; 225: 225-32.
- [63] Belluzzi A, Brignola C, Campieri M et al. Effect of an enteric-coated fish-oil preparation on relapses in Crohn's disease. *N Engl J Med* 1996; 334: 1557-60.
- [64] Maria O'Sullivan, Colm O'Morain. Nutrition in inflammatory bowel disease. *Clinical Gastroenterology* 2006; 561-573.
- [65] Ibrahim A, Aziz M, Hassn A, Mbodjii K, Collasse E et al. Dietary  $\alpha$ -linolenic acid-rich formula reduces adhesion molecules in rats with experimental colitis. *Nutrition* 2012; 28 (7-8): 799-802.
- [66] Marteau P, Boutron-Rualt MC. Nutritional advantages of probiotics and prebiotics. *British J Nutr* 2002; 87: 153S-157S.
- [67] Kelly DG, Fleming CR. Nutritional considerations in inflammatory bowel diseases. *Gastroenterol Clin N Am* 1995; 24: 597-611.
- [68] Dickinson RJ, Ashton MG, Axon ATR et al. Controlled trial of intravenous hyperalimentation and total

- bowel rest as an adjunct to the routine therapy of acute colitis. *Gastroenterology* 1980; 79: 1199-204.
- [69] McIntyre PB, Powell Tuck J, Wood SR et al. Controlled trial of bowel rest in the treatment of severe acute colitis. *Gut* 1986; 27: 481-5.
- [70] Shiloni E, Coronado E, Freund HR. Role of total parenteral nutrition in the treatment of Crohn's disease. *Am J Surg* 1989; 157: 180-5.
- [71] Jones VA, Dickenson RJ, Workman E, Wilson AJ, Freeman AH, Hunter JO. Crohn's disease: maintenance of remission by diet. *Lancet* 1985; 2: 177-80.
- [72] Lochs H, Meryn S, Marosi L et al. Has total bowel rest a beneficial effect in treatment of Crohn's disease? *Clin Nutr* 1983; 2: 61-4.
- [73] Afonso JJ, Rombeau JL. Nutritional care for patients with Crohn's disease. *Hepato-gastroenterology* 1990; 37: 32-41.
- [74] Rombeau JL, Barot LR, Williamson CE et al. Preoperative total parenteral nutrition and surgical *outcome* in patients with inflammatory bowel disease. *Am J Surg* 1982; 143: 139-43.
- [75] Eisenberg HW, Turnbull JRB, Weakley Fl. Hyperalimentation as preparation for surgery in transmural colitis (Crohn's disease). *Dis Colum Rectum* 1974; 17: 469-75.
- [76] Lashner BA, Evans AA, Hanauer SB. Preoperative total parenteral nutrition for bowel resection in Crohn's disease. *Dig Dis Sci* 1989; 34: 741-74.
- [77] Griffiths A, Ohlsson A, Sherman P, et al. Meta-analysis of enteral nutrition as a primary treatment of active Crohn's disease. *Gastroenterology* 1995; 108: 1056-67.
- [78] Fernandez-Berares F, Cabrè E, Esteve-Comas M et al. How effective is enteral nutrition in inducing clinical remission in active Crohn's disease? A meta-analysis of the randomised controlled trials. *JPEN* 1995; 19: 356-62.
- [79] Trallori M, D'Albasio G, Milla M et al. Defined formula diets versus steroids in the treatment of active Crohn's disease. *Scand J Gastroenterol* 1996; 31: 267-72.
- [80] Leiper K, Woolner J, Mullan MMC et al. A randomised controlled trial of high versus low long chain triglyceride whole protein feed in active Crohn's disease. *Gut* 2001; 49: 790-4.
- [81] Hond ED, Hiele M, Peeters M et al. Effect of Long-Term Oral Glutamine Supplements on Small Intestinal Permeability in Patients With Crohn's Disease. *JPEN* 1999; 23: 7-11.
- [82] Akobeng AK, Miller V, Stanton J et al. Double-Blind Randomized Controlled Trial of Glutamine-Enriched Polymeric Diet in Treatment of Active Crohn's Disease. *JPEN* 2000; 30: 78-84.
- [83] Lorenz-Meyer H, Bauer P, Nicolay C et al. Omega-3 Fatty Acid and Low Carbohydrate Diet for Maintenance of Remission in Crohn's Disease. A Randomized Controlled Multicenter Trial. *Scand J Gastroenterol* 1996; 31: 778-85.
- [84] Turner D, Zlotkin SH, Shah PS, Griffiths AM. Omega 3 fatty acids (fish oil) for maintenance of remission in Crohn's disease. *Cochrane Database Syst Rev* 2009 Jan 21; (1).